arXiv:astro-ph/0104447v1 27 Apr 2001

Chandra and *ASCA* X-ray Observations of the Radio Supernova SN1979C IN NGC 4321

A. Ray [1]
Tata Institute of Fundamental Research, Bombay 400 005, India

R. Petre
Lab. of High Energy Astrophysics, NASA/Goddard Space Flight Center, Greenbelt, MD 20771, USA

E. M. Schlegel
Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138, USA

ABSTRACT

We report on the X-ray observation of the radio selected supernova SN1979C carried out with *ASCA* in 1997 December and serendipitously available from a *Chandra* Guaranteed Time Observation in 1999 November. The supernova, of type SN II-Linear (SN $\mathrm{II_L}$), was first observed in the optical and occurred in the weakly barred, almost face on spiral galaxy NGC 4321 (M100). The galaxy, a member of the Virgo S cluster, is at a distance of 17.1 Mpc, and contains at least three other supernovae discovered in this century. The useful exposure time was ∼25 ks for the Solid-State Imaging Spectrometer (SIS), ∼28 ks for the Gas Scintillation Imaging Spectrometer (GIS), and ∼2.5 ks for *Chandra*'s Advanced CCD Imaging Spectrometer (ACIS). No point source was detected at the radio position of SN1979C in a 3' diameter half power response circle in the *ASCA* data. The background and galaxy subtracted SN signal had a 3σ upper limit to the flux of 6.3×10^{-14} ergs $\mathrm{s^{-1}}$ $\mathrm{cm^{-2}}$ in the full *ASCA* SIS band (0.4-10.0 keV) and a 3σ upper limit of $<3\text{-}4\times10^{-14}$ erg $\mathrm{s^{-1}}$ $\mathrm{cm^{-2}}$ in the 2-10 keV band. In the *Chandra* data, a source at the position of SN1979C is marginally detected at energies below 2 keV at a flux consistent with the *ROSAT* HRI detection in 1995. At energies above 2 keV, no source is detected with an upper limit of $\sim3\times10^{-14}$ ergs $\mathrm{s^{-1}}$ $\mathrm{cm^{-2}}$. These measurements give the first ever x-ray flux limit of a Type II_L SN above 2 keV which is an important diagnostic of the *outgoing* shock wave ploughing through the circumstellar medium.

Subject headings: supernovae – supernovae:individual (SN1979C) – galaxies:individual (NGC4321)

1. X-ray and Radio emission from Supernovae

The observation of X-rays provides the most direct view of the region of interaction between the ejecta from a core-collapse supernova (SN) and the circumstellar medium of the progenitor star (see e.g. Chevalier & Fransson 1994). This circumstellar interaction gives rise to copious X-ray emission within days to months of the supernova explosion, in contrast to X-ray emission that occurs in the later supernova remnant (SNR) during the free expansion, adiabatic, or radiative phases. The circumstellar medium is established by the wind-driven mass loss from the progenitor and is heated by the outgoing SN shock wave while the SN ejecta are heated by a reverse shock wave.

---

[1] formerly NAS-NRC Senior Research Associate at NASA/GSFC, Greenbelt, MD 20771; E-mail: akr@tifr.res.in

At the same time, this interaction is expected to produce nonthermal radio radiation; in recent years, several young SNe have been found to be emitting in the radio bands (see Weiler et al. 1998). Relativistic electrons and magnetic fields in the region, which give rise to the radio emission, may be built up by acceleration in the shock wave and field amplification through Rayleigh-Taylor instabilities. The detection of strong radio emission from a SN provides good empirical evidence of circumstellar interaction of ejecta, and such a SN is a probable emitter at X-ray wavelengths as well. Study of the SN X-ray and radio light curves gives an indication of how the shock-front radius changes with time. The latter in turn can be related to the supernova gas radial density profile.

The observations of X-ray emission from the six supernovae discovered prior to 1995 have been reviewed by Schlegel (1995). Currently, there are thirteen known X-ray emitters, most of which are classified as type II SNe on the basis of optical spectra[2]. SN1998bw (a type Ic SN) has been associated with a gamma-ray burst source GRB980425 and is a prototype of the collapsar model involving a rotating massive star (MacFadyen & Woosley 1999). Since the predictions of the models of GRBs regarding afterglows are different on account of different immediate circumsource medium, it is important to detect and monitor the X-ray emission from SNe to understand the processes responsible for the early and long-term electromagnetic emission. In addition, the environments of type II and type Ib SNe are expected to be different due to differing scenarios of mass loss from the progenitor star.

X-ray emission from SNe that turns on shortly after the explosion and remains active for a long time is most likely due to the thermal emission from the hot gas produced in the interaction between the shock wave from the explosion and the circumstellar medium. X-ray emission with a temperature of ≈ 1 keV (which falls into the *ASCA* spectral band) results from a reverse shock that propagates back into the expanding debris. The reverse shock is in turn is developed due to the interaction of the freely expanding supernova ejecta catching up to the decelerating (outgoing) shocked shell (Chevalier & Fransson 1994). The temperature behind the outgoing shock is typically >10 keV and therefore the bulk of this radiation detected by *ASCA* would appear in its higher energy bands. In the circumstellar interaction model, the shocked supernova ejecta is the dominant source of X-rays because of its higher density and lower temperature.

In this paper, we report on an X-ray observation of SN1979C using *ASCA* and an analysis of a *Chandra* observation obtained from the data archive. These observations provide the first constraint on the X-ray flux in the energy band higher than 2 keV for a Type $\mathrm{II_L}$ SN.

2. Observed properties of SN1979C

SN1979C was discovered in the optical by Johnson 1979 near maximum optical light ($m_B^{max} \leq 12$) on April 19, 1979 in the weakly-barred, almost face-on spiral galaxy NGC 4321 (M100). This galaxy is a member of the Virgo S cluster, at a distance of 17.1 Mpc (Freedman et al. 1994). Three other SNe have been observed in this galaxy this century: SN1901B, SN1914A, and SN1959E. The J2000 position of the SN is at $\alpha = 12^h22^m58^s.58$, $\delta = +15^o47^{'}52^s.7$, offset to the southeast from the center of NGC 4321 by ∼100 arc seconds.

SN 1979C has been detected in the radio and X-ray bands and is classified as a type $\mathrm{II_L}$ SN. The radio observation points to a probable explosion date of April 4, 1979 (Weiler et al. 1992). The supernova is thought to have originated from a red supergiant progenitor with $M_{ZAMS} = 17-18\pm3M_\odot$ (Van Dyk et al. 1999). The SN shell expansion velocity near maximum light was $\sim$ 9000 km s^{-1} (see for example, Panagia et al. 1980; Branch et al. 1981; de Vaucouleurs et al. 1981; Fransson et al. 1984; Weiler et al. 1986, 1991; Fesen & Matonick 1993.

SN1979C was observed by the Einstein HRI three times: in 1979 June (∼4 ks), 1979 December (∼13 ks), and 1980 July (∼24 ks). The upper limits for these observations, in the 0.1-4.5

---

[2] The 13 supernovae are: SN1978K (II; Schlegel et al. 1999 and references therein; Ryder et al. 1993), SN1979C (IIL; Immler et al. 1998a), SN1980K (IIL; Canizares, Kriss, & Feigelson 1982), SN1986J (II; Houck et al. 1998), SN1987A (II; Hasinger et al. 1996; Dotani et al. 1987), SN1988Z (IIn; Aretxaga et al. 1999 and references therein), SN1993J (IIb; Zimmermann et al. 1994), SN1994I (Ic; Immler et al. 1998b), SN1994W (IIn; Schlegel 1999), SN1995N (Fox et al. 2000), SN1998bw (Ic; Pian et al. 2000), SN1999em (IIP; Pooley et al. 2001), and SN1999gi (IIP; Schlegel 2001).

keV band and assuming a 5 keV Raymond-Smith plasma, are: June: $L_x \leq 1.8 \times 10^{40}$ erg s^{-1}; December: $L_x \leq 7.6 \times 10^{39}$ erg s^{-1}; and July: $L_x \leq 6.91 \times 10^{39}$ erg s^{-1} at a 3σ confidence level and calculated for a distance of 17.1 Mpc (Palumbo et al. 1981). An analysis of the combined Einstein data, with a total duration ∼41 ks between days 64 and 454 after the explosion, gave an upper limit of 5.9 $\times 10^{39}$erg s^{-1} (Immler et al. 1998a). The foreground column density of neutral hydrogen is $N_H = 2.3 \times 10^{20}$cm^{-2} (using the dust maps of Schlegel, Finkbeiner, & Davis (1998) and the E_{B-V}-N_H relation of Predehl & Schmitt (1995)).

An X-ray observation by the *ROSAT* HRI in 1995 June detected a source coincident with the position of SN1979C for the first time, more than 16 years after the explosion (source H25, Immler et al. 1998a). The *ROSAT* HRI position is $\alpha = 12^h22^m58^s.57$, $\delta = +15^o47^{'}53.5^{"}$ (J2000), within about 2.7$''$ with the position determined by radio interferometry (Penhallow 1980). The measured count rate is 6.9×10^{-4} cps. The corresponding 0.2 - 2.4 keV flux and luminosity are $F_x = 2.9 \times 10^{-14}$ erg cm^{-2} s^{-1} and $L_x = 1.0 \times 10^{39}$ erg s^{-1} respectively, assuming a 5 keV thermal spectrum.

3. Observations

3.1. *ASCA*

NGC 4321 was observed by *ASCA* on December 18-19, 1997. The Solid State Imaging Spectrometers (SIS) had useful exposure times of 24.3 ks (SIS0) and 25.2 ks (SIS1); the Gas Scintillation Imaging Spectrometers (GIS) had exposure times of 27.5 ks (GIS2) and 28.1 ks (GIS3).

The nominal pointing position, $\alpha = 12^h23^m58^s$, $\delta = +15^o51^{'}$, placed the galaxy and the supernova towards the center of the best calibrated SIS chips (chip 1 of SIS0 and chip 3 of SIS1).

A combined SIS0+SIS1 image is shown in Figure 1. A $3^{'}$ diameter half power response circle is shown superposed at the location of SN1979C. The circle approximately matches the size of the D_{25} cicle. Emission associated with the known agglomeration of sources in the central region of NGC 4321 is clearly seen. On the other hand, no point source is evident in the circle. A point source detection task failed to detect SN1979C in either the SIS or GIS image.

Placing an upper limit on the *ASCA* band flux from SN1979C is complicated by contamination from the integrated flux from the sources in the nuclear region of NGC 4321. The center of NGC 4321 is only 100 arc seconds from SN1979C, resulting in a non-negligible overlap of point responses. Furthermore, the emission from the galaxy is clearly broader than that produced by a single unresolved source. Thus determining a count rate from SN1979C requires subtraction both of diffuse background and the contribution of the integrated flux from the other sources in the galaxy.

The background subtracted counts in the defined extraction circles around the galaxy and SN are 178 ±18 counts and 82 ±15 counts respectively. As the overlap between the two circles centered about the galaxy and SN is estimated to be ∼0.4, the background- and galaxy-subtracted SN signal is 11 ±16 counts in the full 0.4 - 10 keV band. The implied 3σ upper limit of the count rate is 2.0 $\times 10^{-3}$ cps. A similar analysis of SIS1 data of 25 ks exposure, combined with the above 24 ks exposure for SIS0, gives a combined upper limit to the count rate of 1.2 $\times 10^{-3}$ cps. The corresponding upper limit to the combined data from GIS2 and GIS3 (28 ks each) gives 8 $\times 10^{-4}$ cps.

The net exposures, total galaxy- and background-subtracted net source counts and their 1 σ errors together with the 3σ upper limits on the SN count rates for SIS (S0 and S1) and GIS (G2 and G3) are given in Table 1.

The above SIS combined upper limit corresponds to a flux limit of 6.3×10^{-14} erg cm^{-2} s^{-1} in the 0.4-10.0 keV band for a thermal bremsstrahlung spectrum with T = 3 keV ($N_H = 2.3 \times 10^{20}$cm^{-2}). For kT = 5 keV, the flux limit is ∼30% higher, ∼8.2×10^{39} erg s^{-1}.

In Table 1 we also list the detected x-ray source fluxes (or the upper limits) for the previous observations obtained with *Einstein* and *ROSAT* along with their respective band-passes and luminosities at different epochs.

4. *Chandra*

The ACIS observation was obtained on 1999 November 6 (observation id = 400; PI=G. Garmire) as part of a survey of low-luminosity galaxy nuclei (Ho et al. 2001). The total exposure time was

2.5 ksec. The nucleus was centered on the back-illuminated (S3) chip; each chip is $8'$ by $8'$ in size, so the exposure, while short, potentially captures emission from SN1979C.

Figure 2 shows the image of the events in the 0.4-2.4 keV band, approximating the *ROSAT* band. The nucleus is clearly visible. Using a circle of radius 3 pixels, which encloses $\sim$92% of the flux, and centered on the position of SN1979C, 21 ± 7 counts are extracted. The corresponding flux and luminosity, for an adopted 3 keV bremsstrahlung spectrum are 4.6×10^{-14} erg s^{-1} cm^{-2} and 1.6×10^{39} erg s^{-1}, respectively. These values are comparable to the values obtained by Immler et al. (1998a) from their *ROSAT* HRI observation. While formally the ACIS measurement shows an increase of a factor $\sim$2 above the *ROSAT* result, that measurement requires confirmation with a longer exposure.

Only an upper limit is possible in the 2-10 keV band; the upper limit is $\sim1.1\times10^{39}$ erg s^{-1} (3σ). This upper limit is similar to the limit on the nucleus reported in Ho et al. (2001). A longer exposure is necessary to establish whether SN1979C is deficient in X-rays above 2 keV.

5. Discussion

The *ROSAT* HRI flux from the measurement of Immler et al. 1998a in the 0.5-2.0 keV band (for the same spectrum and absorption column for SN1979C) is 2×10^{-14}erg cm^{-2} s^{-1} which corresponds to a luminosity of $\sim7\times10^{38}$ ergs s^{-1}. The *ASCA* and *Chandra* flux limits are $\sim3\times10^{-14}$ erg s^{-1} cm^{-2} or a luminosity of 1×10^{39} erg s^{-1}. For the energy bands in common between the two satellites ($\sim$0.5-2.0 keV), *ROSAT* provides the more sensitive measure; above 2 keV, a longer exposure using *Chandra* will clearly reduce the upper limit.

Until now almost all SNe detected in the X-ray were measured in the lower energy band (0.1 - 2.0 keV) of ROSAT. Our reported measurement of SN1979C here gives the flux limit in a harder energy band ($\gtrsim$ 2 keV) for the first time for a Type II$_L$ SN (the only other X-ray detected Type II$_L$ is SN1980K). The outgoing shock wave expanding through circumstellar material creates hotter emission compared to the reverse shock wave. The measurement of SN X-ray fluxes in higher energy bands constitutes an important goal.

Most extragalactic SNe detected so far in the radio are extremely bright compared to galactic supernova remnants. For example, although SN1970G is fainter than about half a dozen radio supernovae (some of which like SN1988Z and SN1986J are nearly 20-30 times as bright at 6 cm), it has a peak spectral luminosity at these wavelengths that is nearly an order of magnitude greater than the brightest galactic supernova remnant, Cassiopeia A (Weiler et al. 1989). Strong radio emission from a supernova provides good empirical evidence of circumstellar interaction and such a supernova is a probable emitter at X-ray wavelengths as well. One expects strong X-ray emission from SN1979C.

To estimate the expected X-ray fluxes from SN1979C on the basis of its radio emission, we scaled its radio flux density (8.3 mJy) with respect to SN1980K (another type II-L SN), as reported by Weiler et al. 1989. The ratio of the respective 6 cm radio flux densities of SN1979C and SN1980K (the latter SN's radio flux is taken from Weiler et al. as 2.6 mJy) is multiplied by the absorption-corrected Einstein X-ray flux of 0.03 μJy at 1 keV as reported by Canizares, Kriss, & Feigelson 1982 for SN1980K, to produce an expected x-ray flux of 1.1 $\times10^{-13}$erg cm^{-2} s^{-1} in the (0.5- 2.0) keV band for SN1979C.

Both the *ROSAT* detection of the x-ray emission as well as the *ASCA* and *Chandra* upper limits lie below this expected value. This lower than expected L_x/L_R may mean that the X-ray and radio emission properties of SNe can be quite variable from one environment to another. In other words, the emissions in the two bands are not directly scalable in a universal way. Or the X-ray emission could be more strongly time variable than the radio emission. Since the X-ray emission from a supernova is found to be primarily from the reverse shock region, the variable outer supernova density profile can affect the X-ray luminosity, but is unlikely to affect the radio luminosity to the same extent.

The detection probability of a SN against a bright galactic background can be limited by a given telescope's half power diameter, especially when a SN fades with time. The higher spatial resolution of *Chandra* improves the detection capability in this regard and is important for their higher energy sensitivity.

6. Acknowledgements

The authors explicitly thank the referee for one of the speediest reports any of us have received. The authors are grateful to the members of the *ASCA* team. One of us (A.R.) was a Senior Research Associate of the National Research Council at NASA/Goddard Space Flight Center during the course of this work and acknowledges support from the NRC and *ASCA* Guest Observer Program. He also thanks the Aspen Center for Physics for hospitality duringa portion of this project. The research of EMS was supported by contract number NAS8-39073 to SAO.

REFERENCES

Aretxaga, I., Benetti, S., Terlevich, R. J., Fabian, A. C., Cappellaro, E., Turatto, M., & Della Valle, M. 1999, MNRAS, 309, 343

Branch, D., Falk, S.W. & McCall, M.L., 1981, ApJ, 244, 780

Canizares, C. R., Kriss, G.A., & Feigelson, E. D. 1982, ApJ 253, L17

Chevalier, R.A., & Fransson, C. 1994 Ap. J., 420, 268

Dotani, T., et al. 1987, Nature, 330, 230

Fesen, R.A. & Matonick, D.M. 1993, ApJ 407, 110

Fox, D. & Lewin, W. 1999, IAU Circular 7318

Fox, D. W., Lewin, W. H. G., Fabian, A., Iwasawa, K., Terlevich, R., Zimmermann, H.-U., Aschenbach, B., Weiler, K., Van Dyk, S., Chevalier, R., Rutledge, R., Inoue, H., & Uno, S. 2000, MNRAS, 319, 1154

Fransson, C., Benvenuti, P. & Wamsteker, W. et al 1984, A & A 132, 1

Freedman, W. et al, 1994, Nature 371, 757

Hasinger, G., Aschenbach, B., & Trümper, J. 1996, A&A, 312, L9

Ho, L. C. et al. 2001, ApJ(Letters), submitted (astro-ph/0102504)

Houck, J. C., Bregman, J. N., Chevalier, R. A., & Tomisaka, K. 1998, ApJ, 493, 431

Immler, S. & Pietsch, W. & Aschenbach, B. 1998, A. & A. 331, 601

Immler, S., Pietsch, W. & Aschenbach, B. 1998, A. & A. 336, L1

Itoh, M., Asai, K., Uno, S., Inoue, H., Nomoto, K., Kumagai, S., Masai, K., & Tanaka, Y. 1997, in X-ray Imaging and Spectroscopy of Cosmic Hot Plasmas, ed. F. Makino & K. Mitsuda, (Tokyo: Universal Academy Press, Inc.), 381

Johnson, G.E. 1979, IAUC 3348

MacFadyen, A. I. & Woosley, S. E. 1999, ApJ, 524, 262

Palumbo, G.G.C. et al 1981 ApJ 247, 484

Panagia, N., Vettolani, G. & Boksenberg, A., 1980 MNRAS 192, 861

Penhallow, W.S. 1980, IAUC 3491

Pian, E. et al. 2000, ApJ, 536, 778

Pooley, D. et al. 2001, ApJ, in press (astro-ph/0103196)

Predehl, P. & Schmitt, J. H. M. M. 1995, A&A, 293, 889

Ryder, S., Staveley-Smith, L., Dopita, M., Petre, R., Colbert, E., Malin, D., & E. M. Schlegel 1993, Ap.J. 416, 167

Schlegel, E. M. 2001, ApJ, submitted

Schlegel, E. M. 1999, ApJ, 527, L85

Schlegel, E. M. 1995, Rep. Prog. Phys., 58, 1375

Schlegel, E. M., Ryder, S., Staveley-Smith, L., Petre, R., Colbert, E. J. M., Dopita, M., & Campbell-Wilson, D. 1999, AJ,

Schlegel, D., Finkbeiner, D., & Davis, M. 1998, ApJ, 500, 525

Van Dyk, S. et al 1999, PASP 111, 313

de Vaucouleurs, G., de Vaucouleurs, A. & Buta, R., et al 1981, PASP 93, 36

Weiler, K.W., Van Dyk, S.D., Montes, M.J., Panagia, N., Sramek, R. A. 1998, ApJ 500, 51

Weiler, K.A., Van Dyk, S., Pringle, J., & Panagia, N. 1992, Ap J. 399, 672

Weiler, K. A., et al. 1991, ApJ, 380, 161

Weiler, K. A. et al. 1989, ApJ, 336, 421

Weiler, K. A., et al. 1986, ApJ, 301, 790

Zimmermann, H.-U. et al. 1994, Nature, 367, 621

This 2-column preprint was prepared with the AAS LaTeX macros v5.0.

Table 1: Observations of SN 1979C in NGC 4321

Instrument	Date	ExpT (ksec)	Band (keV)	Net source Counts	Counts/sec (3σ lim)	Flux
			New Observations			
ASCA SIS	1997 Dec	24 + 25	0.4-10.0	29.6 ± 20	$\leq 1.2 \times 10^{-3}$	$\leq 6.3 \times 10^{-14}$
(S0 + S1)			2.0-10.0	13.3 ± 11	$\leq 6.7 \times 10^{-4}$	$\leq 3.5 \times 10^{-14}$
ASCA GIS	1997 Dec	28 + 28	1.0-10.0	-3.5 ± 16	$\leq 8.4 \times 10^{-4}$	$\leq 4.4 \times 10^{-14}$
(G2 + G3)			2.0-10.0	-12.0 ± 11	$\leq 6.2 \times 10^{-4}$	$\leq 3.2 \times 10^{-14}$
Chandra ACIS	1999 Nov	2.5	0.3-2.5	21 ± 7	$\sim 8 \times 10^{-3}$	$\sim 4.6 \times 10^{-14}$
			2.0-10.0	6.6 ± 7.7	$\leq 2.89 \times 10^{-3}$	$\leq 2.9 \times 10^{-14}$
			Previous Observations			
Einstein HRI	Dec 1980	41	0.1 - 4.5	···	···	$\leq 1.7 \times 10^{-13}$
ROSAT HRI	Jun 1995	42.8	0.1 - 2.4	···	···	2.0×10^{-14}

Note: Flux estimated using absorbed bremsstrahlung spectrum with $kT_{Br} = 3$ keV and column density of $N_H = 3 \times 10^{20}$ cm^{-2}.



Fig. 1.— *ASCA* SIS image of the galaxy NGC4321 (M100) containing the SN1979C. The 3′ (diameter) extraction circle used for the SN1979C is shown centered on its position. SN1979C lies ∼100″ SE of the nucleus.

Fig. 2.— *Chandra* ACIS image of NGC 4321. The nucleus is detected below ∼2 keV; the small circle indicates the location of SN1979C. The extracted counts at SN1979C's location are consistent with the expected flux from the *ROSAT* HRI measurement.